Euclidean Geometry: An Introduction to Mathematical Work Math 3600 Spring 2019

Circles

We have learned quite a bit about basic polygonal shapes, especially triangles, and various species of quadrilaterals. Now we turn our attention to circles. This is the subject of Book III in Euclid's *Elements*. We already have one beautiful theorem about circles, that of Thales, but we'd like to have more.

Read the *Elements* Book III Propositions 1-34. For the following propositions you should work in the axiomatic style of Euclid using I.1-34, III.1-34 and any previously proved results.

9.1 Conjecture. Let *AB* and *AC* be two tangent lines from a point *A* outside a circle. Then *AB* is congruent to *AC*.

Definition. We say that two circles *meet at right angles* if the radii of the two circles to a point of intersection make a right angle.


9.2 Conjecture. Let Γ and Ω be two circles with centers *G* and *O*, respectively. Suppose that these circles meet at two points *A* and *B*. If *GAO* is a right angle, then *GBO* is a right angle.

Definition. A quadrilateral *ABCD* is said to be a *cyclic quadrilateral* if there is a circle Γ such that the four vertices *A*, *B*, *C* and *D* lie on Γ .

9.3 Conjecture. A rectangle is always a cyclic quadrilateral.

9.4 Conjecture (Cyclic Quadrilateral Theorem). Let *A*, *B*, *C* and *D* be four points. The quadrilateral *ABCD* is cyclic if and only if angle *DAC* is congruent to *DBC*.

9.5 Conjecture. Let two circles be tangent at a point *A*. If two lines are drawn through *A* meeting one circle at further points *B* and *C* and meeting the other circle at points *D* and *E*, then *BC* is parallel to *DE*.



Pay special attention to III.16, III.18, III.20, III.21, III.31 and III.32.